

ON THE CASSON INVARIANT CONJECTURE OF NEUMANN–WAHL

ANDRÁS NÉMETHI AND TOMOHIRO OKUMA

ABSTRACT. In the article we prove the Casson Invariant Conjecture of Neumann–Wahl for splice type surface singularities. Namely, for such an isolated complete intersection, whose link is an integral homology sphere, we show that the Casson invariant of the link is one-eighth the signature of the Milnor fiber.

1. INTRODUCTION

Almost twenty years ago, Neumann and Wahl formulated the following conjecture:

Casson Invariant Conjecture. [15] *Let (X, o) be an isolated complete intersection surface singularity whose link Σ is an integral homology 3-sphere. Then the Casson invariant $\lambda(\Sigma)$ of the link is one-eighth the signature of the Milnor fiber of (X, o) .*

The conjecture can be reformulated in terms of the geometric genus p_g of (X, o) as well (see below in 4.1).

The conjecture is true for Brieskorn hypersurface singularities by a result of Fintushel and Stern [8]. This and additivity properties (with respect to splice decomposition) lead to the verification of the conjecture for Brieskorn complete intersections, done independently by Neumann–Wahl [15] and Fukuhara–Matsumoto–Sakamoto [9]. For suspension hypersurface singularities it was verified in [15]. Some iterative generalizations, related with cyclic coverings and using techniques of equivariant Casson invariant and gauge theory, were covered by Collin and Saveliev (cf. [3, 4, 5]).

Recently, Neumann and Wahl have introduced an important family of complete intersection surface singularities, the *splice type singularities*. In [18] they treated the case when the link is an integral homology sphere (the reader may consult in [16, 17, 19] the case of rational homology sphere links too). In [18], they have also verified the above conjecture (by a direct computation of the geometric genus) for special splice type singularities (when the nodes of the splice diagram ‘are in a line’).

The goal of the present article is to verify the conjecture for an arbitrary splice type singularity:

Theorem. *The Casson Invariant Conjecture is true for any splice type singularity with integral homology sphere link.*

2000 *Mathematics Subject Classification.* Primary. 14B05, 14J17, 32S25, 57M27, 57R57. Secondary. 14E15, 32S45, 57M25.

Key words and phrases. normal surface singularities, complete intersection singularities, geometric genus, signature, Neumann–Wahl conjecture, $\mathbb{Z}$ -homology spheres, Casson invariant.

The first author is partially supported by NSF grant DMS-0605323, Marie Curie grant and OTKA grants; the second author by the Grant-in-Aid for Young Scientists (B), The Ministry of Education, Culture, Sports, Science and Technology, Japan.

This theorem can also be judged in the light of the following expectation/conjecture (cf. [18, 19], see also section 6 in [22]): any complete intersection surface singularity with integral homology sphere link is of splice type.

The article is organized as follows. Section 2 is a review of splice type surface singularities. The proof of the theorem is broken into two parts: section 3 contains the analytic part, while the topological/combinatorial part is in the last section 4.

The proof contains an inductive formula for the geometric genus. The inductive step corresponds to the splice decomposition of the link Σ (cf. [7]). But we wish to emphasize that the study of any *analytic* invariant with respect to the splice decomposition is rather delicate since the splice decomposition (gluing), rewritten in the language of plumbing, contains a purely topological step (namely the 0-absorption, cf. [7], §22) which cannot be represented in the world of negative definite plumbing graphs. A new construction of section 4 surmounts this difficulty.

Acknowledgements. The second author thanks Marie Curie Fellowship for Transfer of Knowledge for supporting his visit at the Rényi Institute of Mathematics, Budapest, Hungary, where the work on this paper was done. He is also grateful to the members of the Rényi Institute for the warm hospitality.

2. SPLICE TYPE SURFACE SINGULARITIES (REVIEW)

This section contains a brief introduction of *splice type singularities* in terms of “monomial cycles” (cf. [21, §3], [19, §13]). We always assume that all the links are integral homology spheres.

The splice type singularities, which generalize Brieskorn complete intersections, were introduced by Neumann and Wahl (see [16], [18], [19]). To any resolution (or, negative definite plumbing) graph, associated with a fixed plumbed 3-manifold, one first associates a new weighted tree, called the “splice diagram”. From this one writes down the system of (leading forms of) splice diagram equations. They define a splice type singularity whose link is the original 3-manifold. (In the next presentation the splice diagram is almost ‘hidden’, the interested reader may consult the above references for more details.)

2.1. Basic notations. Let (X, o) be a germ of a normal complex surface singularity and $\pi: \tilde{X} \rightarrow X$ a good resolution, i.e., its exceptional divisor $E := \pi^{-1}(o)$ has only simple normal crossings. Let $\{E_v\}_{v \in \mathcal{V}}$ denote the set of irreducible components of E . The link Σ of the singularity X is an integral homology sphere if and only if E is a tree of rational curves and the intersection matrix $I(E) := (E_v \cdot E_w)_{v, w \in \mathcal{V}}$ is unimodular. Let $(m_{vw})_{v, w \in \mathcal{V}} = -I(E)^{-1}$. Then every m_{vw} is a positive integer. We call an element of a group $\sum_{v \in \mathcal{V}} \mathbb{Z} E_v$ a cycle. For any cycle $D = \sum_{w \in \mathcal{V}} a_w E_w$, we write $m_w(D) = a_w$.

Let $\delta_v = (E - E_v) \cdot E_v$, the number of irreducible components of E intersecting E_v . A curve E_v (or its index v) is called an *end* (resp. a *node*) if $\delta_v = 1$ (resp. $\delta_v \geq 3$). We denote by $\mathcal{E}$ (resp. $\mathcal{N}$) the set of indices of ends (resp. nodes). A connected component of $E - E_v$ is called a *branch* of v .

2.2. Monomial cycle. For any $v \in \mathcal{V}$, let $E_v^* = \sum_{w \in \mathcal{V}} m_{vw} E_w$. Then $E_v^* \cdot E_w = -\delta_{vw}$ for every $w \in \mathcal{V}$, where δ_{vw} denotes the Kronecker delta. An element of a semigroup $\sum_{w \in \mathcal{E}} \mathbb{Z}_{\geq 0} E_w^*$, where $\mathbb{Z}_{\geq 0}$ is the set of nonnegative integers, is called a *monomial cycle*. Let $\mathbb{C}\{z\} := \mathbb{C}\{z_w; w \in \mathcal{E}\}$ be the convergent power series ring in $\#\mathcal{E}$ variables. Then for a monomial cycle $D = \sum_{w \in \mathcal{E}} \alpha_w E_w^*$, we associate a monomial $z(D) := \prod_{w \in \mathcal{E}} z_w^{\alpha_w} \in \mathbb{C}\{z\}$.

2.3. Degree and order associated with v . We fix $v \in \mathcal{V}$. One defines the v -weight of any variable z_w , $w \in \mathcal{E}$, by m_{vw} . Note that $\gcd\{m_{vw}\}_{w \in \mathcal{V}} = 1$. If D is a monomial cycle, then the v -degree of $z(D)$ is equal to $m_v(D)$.

For any $f \in \mathbb{C}\{z\}$ write $f = f_0 + f_1 \in \mathbb{C}\{z\}$, where f_0 is a nonzero quasihomogeneous polynomial with respect to the v -weight and f_1 is a series in monomials of higher v -degrees. Then we call f_0 the v -leading form of f , and denote it by $\text{LF}_v(f)$. We define the v -order of f by $v\text{-ord}(f) := v\text{-deg}(\text{LF}_v(f))$.

2.4. Monomial Condition. We say that E satisfies the monomial condition if for any node v and any branch C of v , there exists a monomial cycle D such that $D - E_v^*$ is an effective cycle supported on C . In this case, $z(D)$ is called an *admissible monomial* belonging to the branch C . Note that a chain C always admits a monomial cycle.

The monomial condition is equivalent to the semigroup and congruence conditions of Neumann-Wahl (see [19, §13]) (although the congruence condition is trivial in our case.)

2.5. Neumann-Wahl system. Assume that the monomial condition is satisfied. For any fixed node v let $C_1, \dots, C_{\delta_v}$ be the branches of v . Suppose $\{m_1, \dots, m_{\delta_v}\}$ is a set of admissible monomials such that m_i belongs to C_i for $i = 1, \dots, \delta_v$. Let $F = (c_{ij})$, $c_{ij} \in \mathbb{C}$, be any $((\delta_v - 2) \times \delta_v)$ -matrix such that all the maximal minors have rank $\delta_v - 2$. We define polynomials $f_1, \dots, f_{\delta_v - 2}$ by

$$\begin{pmatrix} f_1 \\ \vdots \\ f_{\delta_v - 2} \end{pmatrix} = F \begin{pmatrix} m_1 \\ \vdots \\ m_{\delta_v} \end{pmatrix}.$$

We call the set $\{f_1, \dots, f_{\delta_v - 2}\}$ a *Neumann-Wahl system* at v . If we have a Neumann-Wahl system $\mathcal{F}_v$ at every node v then we call the set $\mathcal{F} := \bigcup_{v \in \mathcal{N}} \mathcal{F}_v$ a *Neumann-Wahl system* associated with E . Note that $\#\mathcal{F} = \#\mathcal{E} - 2$.

2.6. Splice diagram equations. Consider a finite set of germs

$$\{f_{vj_v} \mid v \in \mathcal{N}, j_v = 1, \dots, \delta_v - 2\} \subset \mathbb{C}\{z\}.$$

If the set $\{\text{LF}_v(f_{vj_v}) \mid v \in \mathcal{N}, j_v = 1, \dots, \delta_v - 2\}$ is a Neumann-Wahl system associated with E then a system of equations

$$f_{vj_v} = 0, \quad v \in \mathcal{N}, \quad j_v = 1, \dots, \delta_v - 2,$$

is called the *splice diagram equations* and the germs $\{f_{vj_v}\}$ *splice diagram functions*. The germ of singularity defined by the splice diagram equations in $(\mathbb{C}^{\#\mathcal{E}}, o)$ is called a *splice type singularity* (associated with the combinatorics of E).

We note that a splice type singularity is an equisingular deformation of a singularity defined by a Neumann-Wahl system associated with E (cf. [21, (4.)]).

In fact, all the above construction of the splice equations (including the validity of the monomial condition) depends only on the dual graph associated with E , hence only on a fixed plumbing graph of the link Σ .

2.7. Theorem. [18, (2.1)] *Suppose that (Z, o) is a splice type singularity associated with E (or, with a graph of Σ). Then (Z, o) is an isolated complete intersection surface singularity whose link is Σ .*

2.8. End-Curve Condition. We keep the notations of (2.1). We say that $\tilde{X}$ satisfies the *end-curve condition* if for each $w \in \mathcal{E}$ there exists an irreducible curve $H_w \subset \tilde{X}$, not an exceptional curve, and a function $f \in H^0(\mathcal{O}_{\tilde{X}}(-E_w^*))$ such that $\text{div}(f) = E_w^* + H_w$. We call such f an *end-curve function* of E_w . The end-curve condition is equivalent to the fact that for every $w \in \mathcal{E}$ the linear system of $\mathcal{O}_{\tilde{X}}(-E_w^*)$ has no fixed component in E .

Notice that if $\tilde{X}$ satisfies the end-curve condition then so do the minimal good resolution and any resolution obtained from $\tilde{X}$ by blowing-up the singular points of E .

If (X, o) is defined by splice diagram equations, then its resolution $\tilde{X}$ satisfies the end-curve condition (since the coordinate functions serve as end-curve functions). The converse is guaranteed by the following:

2.9. Theorem. [18, (4.1)] *Let $\pi : \tilde{X} \rightarrow X$ be a good resolution of a singularity (X, o) . If $\tilde{X}$ satisfies the end-curve condition, then E satisfies the monomial condition and X is of splice type. In fact, if*

$$\psi : \mathbb{C}\{z_w; w \in \mathcal{E}\} \rightarrow \mathcal{O}_{X,o}$$

is a homomorphism of $\mathbb{C}$ -algebras which maps each $z_w = z(E_w^)$ to an end-curve function of E_w , then ψ is surjective and $\text{Ker } \psi$ is generated by splice diagram functions.*

3. FILTRATION AND p_g -FORMULA

Assume that (X, o) is of splice type, hence its resolution satisfies the end-curve condition. We also assume the presence of a node in E (otherwise the graph is A_n). The main result of this section presents an inductive formula for the geometric genus $p_g(X)$ of X with respect to ‘cutting’ E in two parts (corresponding to a certain ‘splice decomposition’ of Σ). It is convenient to do this cutting near a vertex v which isolates either a ‘star-shaped subgraph’ or a ‘string’. The inductive formula involves a filtration of $\mathcal{O}_{X,o}$ associated with v . It can be proved by repeating the line of arguments of [20] for our slightly more general situation (in [20] v is node); for the convenience of the reader we provide the main points of the proof emphasizing the differences.

Let v_1 be an end-node, i.e., E_{v_1} is an end of the minimal reduced connected cycle containing all nodes of E . Let v_2 be the node which is nearest to v_1 . We fix $v \in \mathcal{V}$ in that branch of v_2 which contains v_1 such that v is not an end. (If v_1 is the only node, let v be any vertex which is not an end.) Let $C_1, \dots, C_{\delta_v}$ denote the branches of E_v .

3.1. Filtration with respect to E_v . The v -order defines a filtration $\{F_n\}$ of $\mathbb{C}\{z\}$ by

$$F_n = \{f \in \mathbb{C}\{z\} \mid v\text{-ord}(f) \geq n\} \quad (n \in \mathbb{Z}_{\geq 0}).$$

This induces a filtration of ideals $I_n \subset \mathcal{O}_{X,o}$ ($n \in \mathbb{Z}_{\geq 0}$) by $I_n := \psi(F_n)$, where $\psi : \mathbb{C}\{z\} \rightarrow \mathcal{O}_{X,o}$ is the natural projection (2.9). Let $\mathcal{G}$ denote the associated graded algebra $\bigoplus_{n \geq 0} \mathcal{G}_n$, where $\mathcal{G}_n = I_n/I_{n+1}$.

3.2. Proposition. (Cf. [19, (2.6)].) *Let $\{f_{wj_w}\} \subset \mathbb{C}\{z\}$ be a set of splice diagram functions as in (2.6), and I the ideal of the polynomial ring $\mathbb{C}[z]$ generated by v -leading forms $\{\text{LF}_v(f_{wj_w})\}$. Then $\mathcal{G} \cong \mathbb{C}[z]/I$ and it is a reduced complete intersection ring.*

Proof. The statement can be reduced to the case when v is a node considered in [19, (2.6)] (or [21, (4.4)]) as follows. Let $D_1, \dots, D_{\delta_{v_1}}$ denote the branches of v_1 ; suppose that D_j is a chain of curves and the variable z_j corresponds to the end of D_j for $1 \leq j \leq \delta_{v_1} - 1$. If E_v is on one of these branches, let D_1 be that branch. Then we may assume that the Neumann-Wahl system satisfies the following properties:

$$\begin{aligned} \text{LF}_w(f_{wj_w}) &= m_{wj_w} + a_{wj_w} m_{w\delta_w-1} + b_{wj_w} m_{w\delta_w}, & w \in \mathcal{N}, \quad 1 \leq j_w \leq \delta_w - 2, \\ m_{v_1 j} &= z_j^{\alpha_j}, & 1 \leq j \leq \delta_{v_1} - 1. \end{aligned}$$

Note that $\text{LF}_v(f_{wj_w}) = \text{LF}_v(\text{LF}_w(f_{wj_w}))$ by [21, (3.8)]; while one gets LF_v of $\text{LF}_w(f_{wj_w})$ by deleting the monomials belonging to a branch containing v . Suppose the admissible monomial $M_{v_1 \delta_{v_1}}$ contains z_t . Then we get that

$$\{\text{LF}_v(f_{wj_w})\} \bigcup \{z_1, z_t\} = \{\text{LF}_{v_1}(f_{wj_w})\} \bigcup \{z_1, z_t\}.$$

In the above-mentioned articles it is shown that the second set is a regular sequence and $\{\text{LF}_{v_1}(f_{wj_w})\} \bigcup \{z_1\}$ defines a curve in $\mathbb{C}^{\#\mathcal{E}}$ smooth off the origin. Since $\{\text{LF}_{v_1}(f_{wj_w})\}$ is also a regular sequence, $\mathcal{G} \cong \mathbb{C}[z]/I$ by [19, (3.3)]. For the reducedness, see the argument at the end of section 3 of [19]. $\square$

3.3. Hilbert series. Let $H(t)$ denote the Hilbert series of the graded ring $\mathcal{G}$, i.e.,

$$H(t) = \sum_{i \geq 0} (\dim \mathcal{G}_i) t^i.$$

From (3.2), by a well-known formula, valid for complete intersections, one has:

3.4. Proposition.

$$H(t) = \prod_{w \in \mathcal{V}} (1 - t^{m_{vw}})^{\delta_w - 2}.$$

From (3.2) one may also read the a -invariant of Goto and Watanabe associated with $\mathcal{G}$ (namely $\sum_{w \in \mathcal{N}, j_w} v\text{-deg}(LF_v(f_{wj_w})) - \sum_{w \in \mathcal{E}} v\text{-deg}(z_w)$, cf. [10, (3.1.4)]):

$$a(\mathcal{G}) = \sum_{w \in \mathcal{V}} (\delta_w - 2)m_{vw}.$$

3.5. Let $\pi': \tilde{X} \rightarrow X'$ be the morphism which contracts each connected component of the divisor $E - E_v$ to a normal point and $\pi_1: X' \rightarrow X$ the natural morphism. Let $E' = \pi'(E_v)$.

3.6. Lemma.

- (1) *There exists $k \in \mathbb{Z}_{>0}$ such that $\mathcal{O}_{\tilde{X}}(-kE_v^*)$ is π -generated.*
- (2) *For any $n \in \mathbb{Z}_{\geq 0}$, $I_n = H^0(\mathcal{O}_{X'}(-nE')) = H^0(\mathcal{O}_{\tilde{X}}([-n\pi'^*E']))$.*
- (3) *$-E'$ is π_1 -ample and π_1 coincides with the filtered blowing up*

$$\text{Projan}_X \left(\bigoplus_{n \geq 0} \pi_{1*} \mathcal{O}_{X'}(-nE') \right) \rightarrow X.$$

- (4) *$H^1(\mathcal{O}_{X'}(-nE')) = 0$ for $n > a(\mathcal{G})$.*

Proof. First we treat (1). Let C be one of the branches of v , and E_u be that component of C which intersects E_v . Fix any end E_w of E in C , and set $A := -\sum_{E_v \subset C} \det(-C) I(C)_{vw}^{-1} E_v$. This is an effective cycle supported by C such that $m_u(A)E_v^* + A$ is a monomial cycle, namely $\det(-C)E_w^*$. Therefore, if we fix two branches C_i ($i = 1, 2$), we can construct two positive integers k_i and effective cycles A_i supported by C_i such that $k_i E_v^* + A_i$ is monomial. Set $k = k_1 k_2$ and $f_i \in H^0(\mathcal{O}_X)$ the image of the monomial $z(kE_v^* + k_j A_i)$, where $i, j = 1, 2$ and $i \neq j$. Then it is easy to see that $f_1, f_2 \in H^0(\mathcal{O}_{\tilde{X}}(-kE_v^*))$ generate $\mathcal{O}_{\tilde{X}}(-kE_v^*)$.

The proof of (2)–(4) is same as the proof of [20, (3.3–3.4)] based on part (1) and filtered ring theory [24]. $\square$

Since $m_{vv}\pi'^*E' = E_v^*$ is integral, from (2) we get for any $m \in \mathbb{Z}_{>0}$

$$(3.1) \quad I_{mm_{vv}} = H^0(\mathcal{O}_{\tilde{X}}(-mE_v^*)).$$

3.7. **p_g -formula.** Let (X_i, x_i) denote the singularity germ obtained by contracting the branch C_i (hence $X_i \subset X'$). Fix $k \in \mathbb{Z}_{>0}$ such that $\pi'_*(kE_v^*)$ is Cartier. Its existence is guaranteed by the proof of (3.6)(1) (namely by the existence of functions f_i).

3.8. **Theorem.** *Set $P(n) = \sum_{i=0}^{n-1} (\dim \mathcal{G}_i)$ for $n \in \mathbb{Z}_{>0}$. Then for any integer $m > a(\mathcal{G})/km_{vv}$*

$$p_g(X, o) = P(mkm_{vv}) - \chi(mkE_v^*) + \sum_{i=1}^{\delta_v} p_g(X_i, x_i).$$

Proof. We write $D = mkE_v^*$ and $D' = mkm_{vv}E'$. Consider the exact sequence

$$0 \rightarrow \mathcal{O}_{\tilde{X}}(-D) \rightarrow \mathcal{O}_{\tilde{X}} \rightarrow \mathcal{O}_D \rightarrow 0.$$

By equation (3.1) one has

$$\dim_{\mathbb{C}} H^0(\mathcal{O}_{\tilde{X}})/H^0(\mathcal{O}_{\tilde{X}}(-D)) = \dim_{\mathbb{C}} I_0/I_{mkm_{vv}} = P(mkm_{vv}).$$

Therefore it suffices to show that

$$(3.2) \quad h^1(\mathcal{O}_{\tilde{X}}(-D)) = \sum_{i=1}^{\delta_v} p_g(X_i, x_i).$$

From the spectral sequence

$$E_2^{i,j} = H^i(R^j \pi'_* \mathcal{O}_{\tilde{X}}(-D)) \Rightarrow H^n(\mathcal{O}_{\tilde{X}}(-D))$$

one gets the exact sequence

$$H^1(\pi'_*\mathcal{O}_{\tilde{X}}(-D)) \longrightarrow H^1(\mathcal{O}_{\tilde{X}}(-D)) \xrightarrow{\alpha} H^0(R^1\pi'_*\mathcal{O}_{\tilde{X}}(-D)) \rightarrow 0.$$

Since $\pi'_*D = D'$ is Cartier and $\pi'^*E' = E_v^*/m_{vv}$, by projection formula and (3.6)(4),

$$H^1(\pi'_*\mathcal{O}_{\tilde{X}}(-D)) = H^1(\pi'_*\mathcal{O}_{\tilde{X}}(-\pi'^*D')) = H^1(\mathcal{O}_{X'}(-D')) = 0,$$

hence α is an isomorphism. Clearly the support of $R^1\pi'_*\mathcal{O}_{\tilde{X}}(-D)$ is in the set $\{x_i\}_i$. Since $D = \pi'^*D'$ and D' is Cartier, we have

$$(R^1\pi'_*\mathcal{O}_{\tilde{X}}(-D))_{x_i} \cong (R^1\pi'_*\mathcal{O}_{\tilde{X}} \otimes \mathcal{O}_{X'}(-D'))_{x_i} \cong (R^1\pi'_*\mathcal{O}_{\tilde{X}})_{x_i}.$$

This proves (3.2). $\square$

3.9. Periodic constants. In the formula of (3.8), $P(mkm_{vv}) - \chi(mkE_v^*)$ is independent of $m \gg 0$. Let us formulate this property. Let $F(t) = \sum_{i \geq 0} a_i t^i$ be a formal power series. Set $P_F(n) = \sum_{i=0}^{n-1} a_i$. Suppose that $P_F(kn)$ is a polynomial function of n for some $k \in \mathbb{Z}_{>0}$. Then for any $k' \in \mathbb{Z}_{>0}$ satisfying this property, the constant terms of $P_F(kn)$ and $P_F(k'n)$ are the same. We call this constant the *periodic constant* of $F(t)$ and denote it by $F|_{pc}$. For example, the formula of Theorem 3.8 is

$$p_g(X, o) = H|_{pc} + \sum_{i=1}^{\delta_v} p_g(X_i, x_i).$$

Note that if two formal power series $F_1(t)$ and $F_2(t)$ have periodic constants then

$$(F_1 + F_2)|_{pc} = F_1|_{pc} + F_2|_{pc}.$$

3.10. Remarks. (1) The message of the above theorem (3.8) is the following. Fix a negative definite resolution graph Γ (or its plumbed 3-manifold Σ) which satisfies the monomial condition. Then there exists a splice type singularity whose link is Σ (via Neumann-Wahl system/construction), and the geometric genus of any such analytic structure is independent of the choice of the Neumann-Wahl system (by 4.3 of [21] and 10.1 of [19]), hence it depends only on the combinatorics of Γ — it will be denoted by $p_g(\Gamma)$. Moreover, the geometric genus of any such analytic structure satisfies the inductive formula (3.8).

(2) The end-curve condition and splice diagram equations are considered for any surface singularity with rational homology sphere links. For this case, and under the assumption that v is a node, a p_g -formula is established in [20].

4. THE PROOF OF THE MAIN THEOREM

4.1. In this section we prove the main Theorem stated in the introduction. For this, we will apply theorem (3.8) in several different situations.

We start with (X, o) , a splice type singularity (whose minimal good resolution $\tilde{X}$ satisfies the end-curve condition), and whose link Σ is an integral homology sphere. Γ will denote the dual resolution graph associated with $\tilde{X}$; $\mathcal{V}$ stays for the set of vertices of Γ .

It is well-known that in the conjecture one may replace the signature σ of the Milnor fiber by the geometric genus p_g of (X, o) via a formula of Durfee and Laufer [6] $\sigma + 8p_g + c(\Gamma) = 0$, where $c(\Gamma) := K_{\tilde{X}}^2 + b_2(\tilde{X})$ (here K denotes the canonical class and b_2 the second Betti number) depends only on the combinatorics of Γ .

If Γ has only one node then Theorem follows from [18, (7.7)]. A different argument runs as follows: The above p_g formula (namely $p_g = H|_{pc}$, cf. 3.9) is valid for any splice type analytic structure (supported by the same topological type), hence we may assume that (X, o) admits a good $\mathbb{C}^*$ -action. Then (X, o) is automatically a Brieskorn complete intersection (cf. [13]), hence we may apply [15] too. Therefore, in the sequel we assume that Γ has at least two nodes.


4.2. In our inductive procedure the following fact is crucial (cf. also with 3.10): for any splice type singularity with (not necessarily minimal) fixed good resolution graph Γ , consider any connected subgraph $\Gamma' \subset \Gamma$. Let E' denote the reduced connected cycle corresponding to Γ' . Then a neighborhood of E' satisfies the end-curve condition (see [20, (2.15)]); hence E' (or Γ') satisfies the monomial condition by (2.9). Therefore if (X', x') denotes the normal surface singularity obtained by contracting E' , then $p_g(X', x')$ is topological computable from Γ' . We write $p_g(X', x') = p_g(\Gamma')$.

4.3. **Starting the induction.** Let $v_1 \in \mathcal{N}$ be an end-node of Γ and $v_2 \in \mathcal{N}$ the node which is nearest to v_1 . Write $\Gamma' \subset \Gamma$ for the branch of v_1 containing v_2 , and denote by w that vertex in Γ' which is connected by v_1 in Γ . Let $H_{\Gamma, v_1}(t)$ be the Hilbert series of the associated graded ring of the filtration with respect to E_{v_1} . Since $p_g(\Gamma'') = 0$ for all the branches Γ'' of v_1 which are chains, by (3.8) we have the following.

4.4. **Proposition.** $p_g(\Gamma) = H_{\Gamma, v_1}|_{pc} + p_g(\Gamma')$.

4.5. One needs to modify this inductive step, since, e.g., in general, Γ' is not unimodular. The next goal is to fit Γ' into a ‘good’ inductive procedure which is compatible with the splice decomposition of Σ (see also (4.11) for more motivation).

Let Δ be the splice diagram of Σ . For the correspondence between Γ and Δ see [7, §22]. We decompose Δ as the splice of two splice diagrams Δ_1 and Δ_2 , where $\Delta_i \ni v_i$.



Let Γ_i be the minimal plumbing graph associated with Δ_i ($i = 1, 2$) (cf. [7]). Next we review how one can recover from Δ_1 and Δ_2 the maximal chain Γ_0 in Γ which connects v_1 and v_2 , and we relate Γ_2 with Γ' . Set $a := \prod_{i=1}^r a_i$ and $d := \prod_{i=1}^s d_i$. Express a/b as a continued fraction $[\alpha_0, \dots, \alpha_m]$, i.e. $a/b = \alpha_0 - 1/(\alpha_1 - 1/(\dots - 1/\alpha_m))$ with $\alpha_i \in \mathbb{Z}_{>0}$ and $\alpha_i \geq 2$ for $i \geq 1$. Similarly, write $d/c = [\beta_0, \dots, \beta_n]$, and consider the chain:

$$\tilde{\Gamma}_0 : \quad v_1 \xleftarrow{-\alpha_m} \dots \xleftarrow{-\alpha_0} \xleftarrow{-\beta_0} \dots \xleftarrow{-\beta_n} v_2$$

This is not a ‘minimal’ graph, and it is equivalent via (topological) plumbing calculus with the (minimal) chain Γ_0 . This calculus runs as follows: by the positivity of the edge determinant $bc > ad$ (cf. [18, §1]), $\alpha_0 = 1$ or $\beta_0 = 1$. Then one successively blows down the vertices whose weight are -1 , and at some moment inevitably a vertex with weight 0 appears; then one makes a 0-absorption by the rule: $\xleftarrow{-e_1} \xrightarrow{0} \xleftarrow{-e_2} \rightarrow \xleftarrow{-e_1 - e_2}$. In this way one gets Γ_0 . On the other hand, the corresponding maximal chain (with determinant c) in Γ_2 has the following form:

$$\Gamma_2 : \quad \xleftarrow{-\beta_1} \dots \xleftarrow{-\beta_n} \boxed{v_2}$$


4.6. **Lemma.** Consider the following resolution graph $\tilde{\Gamma}_2$ which satisfies (1)–(4) below:

$$\tilde{\Gamma}_2 : \quad \boxed{\Gamma^0} \text{ --- } \dots \text{ --- } \boxed{\Gamma^m} \xrightarrow{-1} \boxed{v'_1} \xrightarrow{-1} \boxed{w} \text{ --- } \boxed{\Gamma'}$$

- (1) the subgraph on the left hand side of w is a chain;
- (2) Γ^0 consists of $(\alpha_0 - 1)$ -vertices; the right-end has weight -3 , all the others -2 ;

- (3) Γ^m consists of $(\alpha_m - 2)$ -vertices, all of them with weight -2 ;
- (4) for $0 < i < m$, Γ^i consists of $(\alpha_i - 2)$ -vertices, the right-end has weight -3 and all the others -2 .

Then $\tilde{\Gamma}_2$ is a negative definite plumbing graph whose associated minimal graph is Γ_2 .

Proof. Glue the chain from the left hand side of w with $\tilde{\Gamma}_0$, blow down the (-1) -vertices and use the 0-absorption which identifies  with Γ_2 . $\square$

4.7. Corollary. *There exists a (non-minimal) resolution graph, namely $\tilde{\Gamma}_2$, which represents the splice diagram Δ_2 , it has Γ' as a subgraph, and supports a splice type singularity.*

Proof. We only have to show that $\tilde{\Gamma}_2$ satisfies the monomial condition. Let u be a node of Γ_2 , let C'' a branch of it which contains the distinguished chain involved in the splicing (otherwise the condition is trivial). Let C be the branch of u in Γ which contains C'' . Then there exists an effective cycle A supported on C so that $E_u^* + A$ is monomial in Γ . Then the restriction $A|_{\Gamma'}$ to Γ' has the property that $E_u^*(\Gamma') + A|_{\Gamma'}$ is monomial and has negative intersection with E_w . Then $A|_{\Gamma'}$ can be extended easily on the chain considered in (4.6) to an effective cycle B on $\tilde{\Gamma}_2$ such that $E_u^*(\tilde{\Gamma}_2) + B$ is monomial in $\tilde{\Gamma}_2$. $\square$

Let $H_{\tilde{\Gamma}_2, v'_1}(t)$ be the Hilbert series of associated graded ring of the filtration with respect to $E_{v'_1}$ of $\tilde{\Gamma}_2$. Then, again by (3.8), we have:

4.8. Proposition. $p_g(\Gamma_2) = H_{\tilde{\Gamma}_2, v'_1}|_{pc} + p_g(\Gamma')$.

4.9. Finally we consider the star-shaped graph Γ_1 (which automatically satisfies the monomial condition), hence (again by Theorem 3.8):

4.10. Proposition. $p_g(\Gamma_1) = H_{\Gamma_1, v_1}|_{pc}$.

4.11. As a consequence, from the above three propositions one gets:

$$(4.3) \quad p_g(\Gamma) - p_g(\Gamma_1) - p_g(\Gamma_2) = (H_{\Gamma, v_1} - H_{\tilde{\Gamma}_2, v'_1} - H_{\Gamma_1, v_1})|_{pc}.$$


The main point in this ‘additivity’ formula is the following. Two terms in the expression $\sigma + 8p_g + c(\Gamma) = 0$ (cf. 4.1) satisfy some additivity properties with respect to the splicing. Namely, the additivity $\lambda(\Gamma) = \lambda(\Gamma_1) + \lambda(\Gamma_2)$ (here we identify a plumbing graph with its plumbed 3-manifold) was proved independently by Akbulut-McCarthy, Boyer-Nicas and Fukuhara-Maruyama (according to [2]). Let $b_1(G_i)$ be the first Betti number of the fiber G_i of the fibered knot determined by Γ_i and the splicing data ($i = 1, 2$). Then, by [18, (6.4)] (cf. also with [12, (5.20)]), one has $c(\Gamma) = c(\Gamma_1) + c(\Gamma_2) - 2b_1(G_1)b_2(G_2)$. In particular, in order to run the induction of our proof, we only need to verify

$$p_g(\Gamma) = p_g(\Gamma_1) + p_g(\Gamma_2) + b_1(G_1)b_1(G_2)/4,$$

a fact already noticed in [18, (6.3)]. Therefore the following proposition, together with equation (4.3), implies the theorem of the introduction.

4.12. Proposition. $(H_{\Gamma, v_1} - H_{\tilde{\Gamma}_2, v'_1} - H_{\Gamma_1, v_1})|_{pc} = b_1(G_1)b_1(G_2)/4$.

Proof. Note that in (4.12), each periodic constant independently is a very complicated (Dedekind sum) expression. Nevertheless, their combination provides the simple expression from the right hand side. The proof is based on an explicit computation of the Hilbert series using (3.4). The corresponding entries m_{vw} might be determined from the splice diagrams using identities of [7, §10], or [18, (9.1)]. Note that the ‘unreduced’ splice diagram associated with $\tilde{\Gamma}_2$, having v'_1 as a node, is the following:



We will use the notation of (4.5) and let $\mathcal{V}_2$ be the set of vertices of Γ_2 . Then $m_{v_1 v_1} = ab$ and $m_{v_1 w_i} = ab/a_i$. Consider the integers $p_i := m_{v_1 w_i}/b = a/a_i$ and $q_w := m_{v_1 w}/a$ for all $1 \leq i \leq r$ and $w \in \mathcal{V}_2$ respectively. Finally, set $g(t) = \prod_{w \in \mathcal{V}_2} (1 - t^{q_w})^{\delta_w - 2}$. Then

$$(4.4) \quad H_{\Gamma, v_1}(t) = \frac{g(t^a)(1 - t^{ab})^{r-1}}{\prod_{i=1}^r (1 - t^{p_i b})}.$$

Since for any $w \in \mathcal{V}_2$, $m_{v_1 w}$ in Γ and $m_{v'_1 w}$ in $\tilde{\Gamma}_2$ are the same, one has:

$$(4.5) \quad H_{\tilde{\Gamma}_2, v'_1}(t) = \frac{g(t^a)}{1 - t^b}.$$

Clearly

$$(4.6) \quad H_{\Gamma_1, v_1}(t) = \frac{(1 - t^{ab})^{r-1}}{(1 - t^a) \prod_{i=1}^r (1 - t^{p_i b})}.$$

We define the functions $Q_1(t)$ and $Q_2(t)$ by

$$Q_1(t) = \frac{(1 - t^a)^{r-1}}{\prod_{i=1}^r (1 - t^{p_i})} - \frac{1}{1 - t}, \quad Q_2(t) = g(t) - \frac{1}{1 - t}.$$

Then from (4.4), (4.5) and (4.6) we have

$$(4.7) \quad H_{\Gamma, v_1}(t) - H_{\tilde{\Gamma}_2, v'_1}(t) - H_{\Gamma_1, v_1}(t) = Q_1(t^b)Q_2(t^a) - \frac{1}{(1 - t^a)(1 - t^b)}.$$

On the other hand, consider the characteristic polynomial $P_i(t) = \det(I - th_{1,i})$ (where I denotes the identity matrix) of the monodromy $h_{1,i}: H^1(G_i) \rightarrow H^1(G_i)$ of the fibered knot with Milnor fiber G_i ($i = 1, 2$). Using A'Campo's formula [1], or [7, §11–12] one has

$$P_1(t) = (1 - t) \cdot \frac{(1 - t^a)^{r-1}}{\prod_{i=1}^r (1 - t^{p_i})},$$

$$P_2(t) = (1 - t)g(t).$$

It is known for fibered links in integral homology 3-spheres (or using A'Campo's formula is easy to check) that $P_i(1) = 1$ and the derivative also satisfies $P'_i(1) = b_1(G_i)/2$.

Notice also that $Q_i(t) = (P_i(t) - 1)/(1 - t)$, hence $Q_i(1) = -P'_i(1)$. Therefore, since Q_i 's are polynomials, $(Q_1(t^b)Q_2(t^a))|_{pc} = Q_1(1)Q_2(1) = b_1(G_1)b_1(G_2)/4$. We end the proof with the remark that $(1 - t^a)^{-1}(1 - t^b)^{-1}|_{pc} = 0$ (since $\gcd(a, b) = 1$). $\square$

REFERENCES

- [1] A'Campo, N.: La fonction zêta d'une monodromie, *Comment. Math. Helv.*, **50**, 233-248, 1975.
- [2] Boyer, S. and Nicas, A.: Varieties of group representations and Casson's invariant for rational homology 3-spheres, *Trans. AMS* **322** (2), 507-522, 1990.
- [3] Collin, O.: Equivariant Casson invariant for knots and the Neumann-Wahl formula, *Osaka J. Math.*, **37** (2000), 57-71.
- [4] Collin, O. and Saveliev, N.: A geometric proof of the Fintushel-Stern formula, *Adv. in Math.*, **147**(1999), 304-314.
- [5] Collin, O. and Saveliev, N.: Equivariant Casson invariants via gauge theory, *J. reine Angew. Math.*, **541** (2001), 143-169.
- [6] Durfee, A.: The Signature of Smoothings of Complex Surface Singularities, *Math. Ann.*, **232**, 85-98, 1978.
- [7] Eisenbud, D. and Neumann, W.: *Three-Dimensional Link Theory and Invariants of Plane Curve Singularities*, Ann. of Math. Studies **110**, Princeton University Press, 1985.
- [8] Fintushel, R. and Stern, R.J.: Instanton homology of Seifert fibered homology 3-spheres, *Proc. London Math. Soc.*, (3) **61**, 109-137, 1991.
- [9] Fukuhara, S.; Matsumoto, Y. and Sakamoto, K.: Casson's invariant of Seifert homology 3-spheres *Math. Ann.* **287** (1990), no. 2, 275-285.
- [10] Goto, S. and Watanabe, K.-i.: On graded rings, *J. Math. Soc. Japan* **30** (1978), No 2, 179-213.
- [11] Laufer, H.B.: On μ for surface singularities, *Proceedings of Symposia in Pure Math.*, **30**, 45-49, 1977.

- [12] Némethi, A. and Nicolaescu, L.I.: Seiberg-Witten invariants and surface singularities, *Geometry and Topology*, **6** (2002), 269-328.
- [13] Neumann, W.: Abelian covers of quasihomogeneous surface singularities, *Proc. of Symposia in Pure Mathematics*, vol. **40**, Part 2, 233-244.
- [14] Neumann, W.D.: A calculus for plumbing applied to the topology of complex surface singularities and degenerating complex curves. *Transactions of the AMS* **268** (1981), no. 2, 299-344.
- [15] Neumann, W. and Wahl, J.: Casson invariant of links of singularities, *Comment. Math. Helv.* **65** (1991), 58-78.
- [16] Neumann, W. and Wahl, J.: Universal abelian covers of surface singularities, *Trends in singularities*, 181-190, *trends Math.* Birkhäuser, Basel, 2002.
- [17] Neumann, W. and Wahl, J.: Universal abelian covers of quotient-cusps, *Math. Ann.* **326** (2003), no. 1, 75-93.
- [18] Neumann, W. and Wahl, J.: Complex surface singularities with integral homology sphere links, *Geometry and Topology*, **9** (2005), 757-811.
- [19] Neumann, W. and Wahl, J.: Complete intersection singularities of splice type as universal abelian covers, *Geometry and Topology*, **9** (2005), 699-755.
- [20] Okuma, T: The geometric genus of splice-quotient singularities, [arXiv:math.AG/0610464](https://arxiv.org/abs/math/0610464).
- [21] Okuma, T: Universal abelian covers of certain surface singularities, *Math. Ann.* **334** (2006), 753-773.
- [22] Stevens, J.: Universal abelian covers of superisolated singularities, [arXiv:math.AG/0601669](https://arxiv.org/abs/math/0601669).
- [23] Tomari, M.: A p_g -formula and elliptic singularities, *Publ. R. I. M. S. Kyoto University*, **21**, 297-354, 1985.
- [24] Tomari, Masataka and Watanabe, Kei-ichi: Filtered rings, Filtered Blowing-Ups, Normal Two-Dimensional Singularities with “Star-Shaped” Resolution, *Publ. R. I. M. S.*, Kyoto University, **25**, 681-740, 1989.

DEPARTMENT OF MATHEMATICS, OHIO STATE UNIVERSITY, COLUMBUS, OH 43210; AND

RÉNYI INSTITUTE OF MATHEMATICS, BUDAPEST, HUNGARY

E-mail address: nemethi@math.ohio-state.edu; nemethi@renyi.hu

DEPARTMENT OF EDUCATION, YAMAGATA UNIVERSITY, YAMAGATA 990-8560, JAPAN.

E-mail address: okuma@e.yamagata-u.ac.jp